

This is a repository copy of *Proximity and High Density of Convenience Stores Was Associated With Obesity in Children of a Rural Community of Mexico: Using a Geographic Information System Approach*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/177482/>

Version: Accepted Version

Article:

Zavala, Gerardo A orcid.org/0000-0002-9825-8725, Yessica, Tenorio-Palos, Maiza, Campos-Ponce et al. (5 more authors) (2021) Proximity and High Density of Convenience Stores Was Associated With Obesity in Children of a Rural Community of Mexico: Using a Geographic Information System Approach. Food and nutrition bulletin. 3795721211033146. ISSN 1564-8265

<https://doi.org/10.1177/03795721211033146>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Proximity and high density of convenience stores was associated with obesity in children of a rural community of Mexico; using a Geographic Information System (GIS) approach

Zavala, Gerardo Antonio^{1,2,3†}; Tenorio-Palos, Yessica^{1†}; Campos-Ponce, Maiza²; Elton-Puente Juana Elizabeth²; López-González Carlos Alberto², Doak Colleen M⁴; Rosado Jorge L¹; García Olga P^{1*}

¹Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, Querétaro, Mexico

²Faculty of Earth and Life Sciences, VU Amsterdam University, Amsterdam, Netherlands

³Faculty of Health Sciences, University of York, York, United Kingdom

⁴Public Health, St Ambrose University, Iowa, U.S.A.

[†]These authors contributed equally in the manuscript

***Corresponding Author:** Olga P. García

Mailing Address: School of Natural Sciences, Universidad Autónoma de Querétaro, Av Ciencias SN Juriquilla, Querétaro, México 76230

Telephone: +52-442-192-1200 ext 5323; 5351

Email: olga.garcia@uaq.mx

Abstract

Background: Food environment (FE) has been linked to obesity in urban areas, but there is limited information in rural areas, particularly in developing countries, where prevalence of obesity is high. **Objective:** To determine the association between FE and childhood obesity using Geographic Information Systems (GIS). **Methods:** A total of 218 children (8–10 y) participated in a cross-sectional study. Weight, height and body fat were measured. Geolocation of convenience stores (CS) and participants' households was collected and the amount of processed food (PF) in the stores was measured. The proximity to the nearest CS and the number of CS within a 250m buffer from each participant's household was calculated using GIS. Linear regression models between obesity measurements and food environment were performed. **Results:** The combined prevalence of overweight and obesity was 32%. A total of 91% of the children had access to a CS within 250 m. On average, 48% of the shelf space of the CS were occupied with PF. A positive association between the density of CS with body fat % ($\beta=0.145$; 95CI: 0.048, 0.241, $p=0.004$), abdominal fat % ($\beta=0.206$; 95%CI: 0.048, 0.241, $p=0.003$) and BMI-for-age z-score ($\beta=0.028$; 95%CI: 0.005, 0.062, $p=0.005$) was found. Living closer to CS was associated with increases in body fat% ($\beta=-0.009$; 95CI: -0.017, -0.001, $p=0.025$), abdominal fat % ($\beta=-0.012$; 95%CI: -0.023, -0.001, $p=0.033$) and BMI-for-age z-score ($\beta=-0.002$, 95%CI: -0.004, -0.001, $p=0.003$). **Conclusion:** In a rural community in Mexico, a high density and low proximity to CS is associated with obesity in school-aged children.

Key Words: Childhood obesity, food environment, GIS, rural areas

Introduction

Almost thirty percent of Mexican children are either obese or overweight. According to nutrition and health surveys, prevalence of childhood obesity has stabilized in urban areas while it still rising in rural areas of Mexico ^{1,2}. Overweight and obese children have an increased risk of oxidative stress, inflammation and insulin resistance and they are more likely to remain obese throughout their life ³. Obesity is the result of the balance between energy intake and expenditure and a complex interaction with socioeconomic, physiological and genetic factors⁴. In addition, new insight has proved that food environment, which has been defined as the entire environment that influence food selection and food intake⁵, plays a key role in the etiology of obesity^{6,7}. The food environment includes food stores and food service establishments, such as convenience stores (CS), supermarkets, local markets, restaurants, fast food stands, among others ^{8,9}. In the developing world, it is also possible to find small CS or neighborhood CS, and these stores represent an important economic model for the retailing business¹⁰. In Mexico and some parts of Central America, these small CS are called “tienditas” or “misceláneas”, are usually family-owned, and are mainly run by the women in the households¹¹. By definition, “tienditas” are convenience stores that provide rapid access to essential products such as food staples (tortillas, rice, beans, eggs, bread, sugar, vegetables, canned foods), toiletries (soap, detergents, toothpaste, toilet paper), processed foods (candy, chips, cookies), and some sell other items such as school supplies¹¹.

Studies addressing the association between food environment and obesity have shown contrasting results among ethnicities, age groups, areas (urban vs rural) and countries ¹²⁻¹⁸. For instance, proximity to supermarkets in Indiana, USA, has been

associated with a healthier diet and negatively associated with overweight and obesity in children in adolescents (3 – 18 y) ¹⁴⁻¹⁶. In contrast, availability of CS within a 0.25 mile (400m) buffer was associated with a greater risk of overweight and obesity in 6-7 year old girls living in Northern California ¹⁷. In addition, it has been observed that income and ethnicity are not only associated with the location, size and number of stores present in the food environment, but also with the selection of available food¹⁸. It is important to consider that most of the studies on this topic have come from developed countries, in urban settings, and have been done in adult populations. Interestingly, even though Mexico has one of the highest rates of obesity worldwide, there is limited information concerning the association between food environment and obesity, and the available information is from urban areas. For instance, one study in two urban settings found that the number of mobile food vendors around schools was positively related with children's BMI¹⁹. In rural settings from developing countries, most families get their food from CS and it is common for children to play a role in household food purchasing ¹¹.

Using Geographic Information Systems (GIS) provides a reliable and objective measurement on food environment ²⁰. A large list of proxies have been used to evaluate food environment, such as the stores within a given radius, the proximity to the nearest food store and the shelf-length of highly processed food and non-processed foods ²¹. A major challenge is to identify proxies that better represent food environment and that consider social, racial, physical and age-related factors in a community or an individual. ^{20,22}.

Given the high rates of overweight and obesity in Mexico and the large proportion of the population living in rural communities, the main objective of this cross-sectional

study is to assess the association between obesity and food environment in children of rural Mexico.

Methods

Population and study design

A cross sectional study was conducted among 218 children (8 -10 years) from a rural community named Santa Cruz in the municipality of El Marques within the state of Queretaro, Mexico. The community has a population of 3902, the ethnicity is predominantly mestizo (mixed-race individual), and agriculture is the main economic activity. The nearest supermarket is in the city of Queretaro (nearest metropolitan center), 35 km away from El Marques, so people from the community usually buy their food from small CS ("tienditas" or "misceláneas"). Children's information was collected from August 2016 until May 2017.

Children were recruited from the elementary school in the community. Parents were asked to attend a meeting where the study procedures were explained. Those that agreed to participate, signed consent forms and were included in the study. Also, children provided their assent to participate. Children who were under a prescribed physical activity or diet regimen were excluded from the study. This study was conducted according to the guidelines of the Declaration of Helsinki and all procedures involving human subjects were approved by the Bioethics Committee of the Universidad Autónoma de Querétaro (UAQ).

Demographic questionnaire

A demographic questionnaire was answered by the caretaker. The questionnaire included information related to their living conditions (number of rooms in the house, inside running water, home appliances, among others), main mode of transportation, family income and caregiver's level of education. The information in the questionnaire was used to determine the socioeconomic level of the participants' households.

Physical activity

Physical activity was estimated using a validated questionnaire specifically for Mexican populations²³. According to the compendium of physical activities of Ainsworth²⁴, the intensities of the physical activities were categorized in three according to the number of the metabolic equivalent of task (METs): light (<3 METs), moderate (3–6 Mets) and vigorous activity (>6 Mets). The resulting hours of each type were divided into days of the week to obtain daily hours practiced of each type of activity.

Anthropometry and body composition

Children were transported with their mothers from the community to the Nutrition Clinic at the UAQ. Weight, height and waist circumference were measured twice with a precision of 0.1 g or 0.1 cm respectively, in all children by trained personnel following the World Health Organization (WHO) procedures²⁵. Children were weighed barefoot and wearing light clothes, using a calibrated digital scale (SECA, mod 813 Hamburg, Germany). Height was measured using a stadiometer (SECA, mod 206 Hamburg, Germany). Nutritional status was calculated based on the WHO criteria of BMI-for-age for children aged 5-19 years using the Anthroplus software (Geneva: WHO, 2009). Underweight was defined as two z-scores below the WHO reference median, overweight

as one standard deviation above the WHO reference median and obese as two standard deviations above the reference median of the BMI-for-age z-score (BMIz)²⁶. Whole body composition was also measured by a certified technician using Dual-energy X-ray absorptiometry (DXA) (Hologic Mod Explorer, 4500 C/W QDR, INC 35 Crosby Drive, Bedford, MA 01730, USA). Body fat percent and body fat content (Kg) were recorded from the values provided by the DXA. Abdominal fat percent and abdominal fat content (Kg) were estimated following procedures described by Hill *et al*²⁷. High body fat for girls was considered above 30% and above 25% for boys^{28,29}.

Geographical data

The geolocation of each participant's household, of all the CS available in the community, and of the school in the community were obtained using "Open Data Kit tool" that was preinstalled in android OS devices equipped with a Global Positioning System (GPS) chip. We identified through a community audit, two types of stores in the community. The first stores that were identified were the "tienditas" or "misceláneas", typical family owned found in Mexico. The second type of food stores found were "food venues", which are local or ambulant small stalls that usually sell 1 or 2 products, such as "aguas frescas" (sugar sweetened fruit drinks), homemade fried chips, candies, among others. For the purpose of this study, we analyzed only CS, since this is where the inhabitants acquire most of their food³⁰. The community audit was done by foot and data collection was obtained from August 2016 until May 2017.

In Store Availability

Using a measuring tape, a two person team measured the shelf-length (m) of Processed Foods (PF) and Non or Minimally Processed Foods (NMPF) to the nearest cm

³¹. The PF were defined as those foods processed to help preserve and enhance nutrients and freshness, foods that combined ingredients (such as flavor, sweeteners, spices and preservatives) for safety and taste, and ready-to-eat foods with minimal or no needed preparation³². These PF included foods such as sugar sweetened beverages, salty snacks (chips, popcorn), cookies, pastries, candies (chocolates, hard candy and gum), ice cream, dairy products (except milk), instant soup, doughnuts, box bread, ready-to-eat cereals, ham, sausages, and canned products. The NMPF included fresh fruits and vegetables, unprocessed chicken, beef or pork, eggs, milk, maize tortillas, beans, rice, and water, among others.

Food Store Access

A GIS database was constructed geocoding all the CS and the participants' households using ArcMap 10 (Redlands, CA) (Figure 1). To determine the best density (number of stores/area) proxy for the food environment, four circular buffers of 150 m, 200 m, 250 m and 300 m radius were defined around each household. The WGS1984 coordinate system with a fuller projection was used to geocode addresses and calculate distances and buffers. The number of stores and the shelf length of each type of food (PF and NMPF) within each buffer was quantified. The length of the buffers was determined based on key GIS data collection information and the characteristics of the community, such as the extension, population and number of stores³³. The proximity was assessed using the Euclidean distance (straight-line distance), to evaluate distance between each participant's household and the nearest convenience store.

Statistical analysis

A descriptive analysis of the demographic variables was performed. As a first step, we assessed which density measure best correlated with obesity. For this purpose, a set of linear regression models between the different measurements of obesity (i.e. BMI, abdominal and total body fat percent) with the number of stores in 150 m, 200 m, 250 m and 300 m buffers were performed. The 250 m buffer was selected to calculate the density of CS because it had the highest coefficient with body fat ($b=0.16$), abdominal fat ($b=0.22$) and BMI ($b=0.03$).

Normality of the independent variables was confirmed by visual inspection and the Shapiro-Wilk test. Linearity of the association was confirmed visually inspecting the data with a LOWESS graph and by plotting the regression residuals. Also, data analysis of all variables were checked for outliers.

We evaluated the relationship between in-store food availability (i.e. total shelf length of PF and NPF) and the different measurements of obesity. We performed a linear model assessing the association between obesity and food environment (i.e. density and proximity of CS), while controlling for the different components of food environment and possible confounders (i.e. sex, age, caregiver's educational level, physical activity and proximity to school). All statistical analyses were performed by SPSS v23.0 (SPSS Chicago, IL, USA).

To evaluate if "other food venues" have an influence on the measurements of obesity, we carried out sensitivity analysis. We calculated the density and proximity, this time including "other food venues" together with the CS in the linear models (i.e. any food venue or convenience store in a 250m buffer or the shortest distance to any "other food venues" or CS).

Results

The general characteristics of the children who participated in the study are shown in Table 1. Of the total population, 55% percent of the studied children were girls, and 52% of the children had an elevated percentage of body fat. The main mode of transportation inside the community in all participants was walking. More than 90% of the households had low SES, more than 29% of the caregivers did not complete elementary school and 91% of the households had access to at least 1 CS within 250 m for their households.

A total of 58 food venues were found in the whole community: 40 CS and 18 other food venues such as two “tortillerías” (stores that only sell tortillas), one pastry shop, two butcheries and one poultry shop, one fast food restaurant and 11 street vendors that sold different kinds of food such as “tamales”, corncob, peanuts, and salty deep-fried “churros”. Sixty-two percent (± 16 SD) of the shelf-space in the CS contains PF, which represents an average of 16.73 m (± 9.3 SD) of the shelf-space (Table 2).

As summarized in Table 3, children with high body fat percent had access to more CS compared to children with normal body fat percent. No differences were observed in proximity to CS in children depending on their body fat percent.

As observed in Table 4, when adjusting for confounding (sex, age, caregiver’s educational level, physical activity and proximity to school), the density of CS was positively associated with all the measurements of obesity (body fat %, abdominal fat % and BMIz). The proximity to CS was negatively associated with all the measurements of obesity, an increase of 100 meters in the proximity to the nearest convenience store was

associated with lower body fat -0.9% (95%CI -1.7% and -0.1%) abdominal fat -1.12% (95%CI -2.3% and -0.1%) and BMI-for-age-z-score -0.2 (95%CI -0.4% and -0.1%). Also, the shelf-meters of both PF and NMPF were positively associated with higher BMIz. However, no differences were observed between groups when analyzing the PF/NMPF ratio.

The sensitivity analysis showed no variations in the coefficient or the direction of the associations between density and proximity with the measurements of obesity after including “other food venues”.

Discussion

In the present study, the food environment of a rural community of Mexico was evaluated. To the best of our knowledge, the results show for the first time the relationship between obesity and measures of food environment in a rural setting in Mexico, where no supermarkets are available, and where its habitants are compelled to buy their food in small CS. In this community, a high density and low proximity to CS was associated with obesity in children, and they were exposed on average to 11.5 CS within 250 m from their household.

The food environment of this community differs importantly from other studies in different countries. The main reason of these differences is that most of the studies that have evaluated food environment and its relationship with obesity, have been done in high income countries, in both urban and rural communities. There is limited research of this type in rural communities from lower-medium income countries, such as Mexico. This has to be taken into consideration when comparing the results from a rural community in

Mexico to those from other settings in high income countries with different contexts. For instance, in a community in Texas, near the border between Mexico and the United states, three CS could be found within a 1600 m buffer of the participants' households⁷. In Denmark, in metropolitan areas only 54% of the population had one or more CS within a 250 m buffer and in non-metropolitan areas only 22% of the population had one or more CS within a 250 m buffer ³⁴. In the city of New Orleans in the US, 38% of the studied households had one store within 1600 m buffer and 62% had the nearest store farther than 1600 m buffer³⁵. In contrast, in the present study, 91% of the children living in the studied community had access to at least 1 CS in a 250 m buffer. Thus, when comparing with other populations and areas in developed countries, the children of this rural community had access to a higher number of CS in their environment.

Higher density and proximity of food stores was associated with a higher BMI, and with higher body and abdominal fat in 8-10 year children living in a rural setting. Most of the studies that have evaluated the association of food environment and obesity have been done in adults and in urban areas, and only a few studies have been done in rural settings and in children^{7,19,36-39}. Results of studies in adults in both urban and rural area are similar to the results of the present study. For instance, in urban settings in the US, proximity, and frequent use of CS, are associated with higher BMI in adults and households^{37,40,41}. Food swamps with a higher density of establishments selling high calorie food and lower density of establishments selling healthier food are predictors of adult obesity, particularly in counties with income inequality and low mobility⁴².

Similarly, studies in rural communities have found that limited access to healthy foods is one of the main contributors to a poor diet and obesity⁴³. Studies that have

evaluated the role that the food environment has on childhood obesity have produced inconsistent result^{44,45}. In Canada, 58% of children had access to one CS within 800 m walking, and proximity and density was not associated with obesity⁴⁶, while children living near a CS in Guam were most likely to have higher BMI z-score compared to those that lived farther away⁴⁷. Similarly, the presence of one or more CS around participants households (a block around or in a 1600 m buffer) in different cities in the US was positively associated with obesity in both children and adolescents^{6,7}. In addition, the presence of CS near schools has also been associated with obesity in different countries^{19,38,48}. In Mexico, for instance, the number of mobile food venues around both, public and private schools, was positively associated with obesity¹⁹). Thus, the evidence suggests that the density of CS near children's households and schools is strongly associated with higher BMI.

The association between the density of food venues and obesity in the population studied may be explained by different social and psychological factors. It is more likely that children living close to a higher density of stores can find a store in their way to their destination (i.e. school, family member house, while playing in the street) and therefore have increased access to CS and food products compared with children living far away from CS^{49,50}. Also, children sharing a higher density of stores are exposed to visual cues more often, both outside and inside the stores, where advertisement as promotions, free toys, and colorful labels of their food products may be found^{51,52}.

In the present study, shelf-space dedicated to PF, which tend to be more energy dense, was higher than NMPF, and the shelf-space of both was associated with obesity in children. However, no association was observed of PF/NMPF with any measure of

obesity. These results suggest that the density and proximity of CS are better predictors of obesity than the proportion of PF/NMPF. To our knowledge, no studies have evaluated the relationship between shelf-space of specific foods and obesity in Mexico, but there is evidence in developed countries. Studies in both Australia and the United States have shown that shelf-space dedicated to energy dense food and less healthy foods is greater in neighborhoods with low SES^{39,53}. Rural stores in Australia had lower shelf space of fruits and vegetables and higher shelf space of soft drinks compared to urban stores, and rural areas in this country have high prevalence of overweight and obesity⁵⁴.

Some limitations of the study need to be addressed. The cross-sectional design of the study does not allow to draw conclusions about causality. Even though associations between food environment and obesity were observed, these associations were small. Since obesity is a multifactorial disease, other factors, such as genetics, may be more important. In addition, there may be behaviors that were not measured in the study, such as the routes the children take to visit family members that may influence their exposure to the food environment of the community. Another limitation is the small sample size of the participants. Also, no information of how often children actually shop in the CS is available, or what type of food they actually purchase. Despite adjusting for proximity to the school, children shared similar food environments near their houses and their school, and thus, the assumption of independence of residuals might be violated. Physical activity was measured using self-reported data and this may be subject to recall bias.

The main strength of the study was the assessment of all the CS available to the community, providing better estimates for the associations found. Another strength of the study was using DXA to measure body composition as an indicator of obesity. Also, the

present study is one of the few addressing the relationship between “tienditas” and childhood obesity in rural communities, where supermarkets are not available.

Future studies in rural communities should include information regarding how often the children buy food in the CS, what type of food they purchase, and how children move within the community, around their households and their school.

Conclusion

Georeferenced locations provide a useful approach to assess the relationship between food environment and obesity by modelling spatial accessibility (density, proximity to CS and exposure to processed foods). The results show that food environment near participants’ households is associated with obesity in 8-10 year old children living in a rural community in Mexico. In a country where prevalence of childhood obesity is one of the highest in the world, understanding the food environment will contribute to shape the environment to promote and support healthy eating and thus, prevent childhood obesity.

Author Contributions

OPG, MCP and GAZ conceived and designed the study. YTP and GAZ carried out the field work, YTP, JLR, JEEP, CALG, CMD and GAZ analyzed and interpreted data, and OPG, CALG and JLR gave important intellectual advice. OPGO, GAZ and YTP drafted the manuscript. All authors were involved in revising the paper and had final approval of the submitted and published versions.

Acknowledgements

The authors will like to acknowledge the children and their parents who gave their consent and support to the study, Dolores Ronquillo, Lorena Oropeza, Paulina Estrella, and Gabriela Curiel who participated in the field work and data management.

Conflict of interest

The Authors declare that there is no conflict of interest.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The project was partially supported by the research network Obesity Treatment and Prevention, financed by CONACYT, Mexico.

References

1. Hernández-Cordero S, Cuevas-Nasu L, Morales-Ruán M, Humarán IM-G, Ávila-Arcos M, Rivera-Dommarco J. Overweight and obesity in Mexican children and adolescents during the last 25 years. *Nutrition & diabetes*. 2017;7(3):e247.
2. Shamah-Levy T, Cuevas-Nasu L, Rivera-Dommarco J, Hernández-Ávila M. Encuesta Nacional de Nutrición y Salud de Medio Camino 2016 (ENSANUT MC 2016). *Informe final de resultados Recuperado de [https://www.insp.mx/ensanut/medio-camino-16/html/\[Links\]](https://www.insp.mx/ensanut/medio-camino-16/html/[Links])*. 2016.
3. Laura Anca P, Bogdana V, Olivia T, Horia V, Dumitru O, Leon Z. The Relations Between Immunity, Oxidative Stress and Inflammation Markers, in Childhood Obesity. *Free radical biology & medicine*. 2014;75 Suppl 1:S44-45.
4. Friedman JM. Obesity: Causes and control of excess body fat. *Nature*. 2009;459(7245):340-342.
5. Timperio A, Crawford D, Leech RM, Lamb KE, Ball K. Patterning of neighbourhood food outlets and longitudinal associations with children's eating behaviours. *Preventive medicine*. 2018;111:248-253.
6. Galvez MP, Hong L, Choi E, Liao L, Godbold J, Brenner B. Childhood obesity and neighborhood food-store availability in an inner-city community. *Academic pediatrics*. 2009;9(5):339-343.
7. Laska MN, Hearst MO, Forsyth A, Pasch KE, Lytle L. Neighbourhood food environments: are they associated with adolescent dietary intake, food purchases and weight status? *Public health nutrition*. 2010;13(11):1757-1763.
8. Wang MC, Gonzalez AA, Ritchie LD, Winkleby MA. The neighborhood food environment: sources of historical data on retail food stores. *International Journal of Behavioral Nutrition and Physical Activity*. 2006;3(1):1.
9. Leung CW, Laraia BA, Kelly M, et al. The Influence of Neighborhood Food Stores on Change in Young Girls' Body Mass Index.

- 367 10. Liese AD, Weis KE, Pluto D, Smith E, Lawson A. Food store types, availability, and cost of foods in
368 a rural environment. *Journal of the American Dietetic Association*. 2007;107(11):1916-1923.
- 369 11. Pisani MJ, Yoskowitz DW. A Sstudy of small neighborhood "tienditas" in Central America. *Latin*
370 *American Research Review*. 2012:116-138.
- 371 12. Morland KB, Evenson KR. Obesity prevalence and the local food environment. *Health & place*.
372 2009;15(2):491-495.
- 373 13. Zenk SN, Powell LM. US secondary schools and food outlets. *Health & place*. 2008;14(2):336-346.
- 374 14. Michimi A, Wimberly MC. Associations of supermarket accessibility with obesity and fruit and
375 vegetable consumption in the conterminous United States. *International Journal of Health*
376 *Geographics*. 2010;9(1):1.
- 377 15. Inagami S, Cohen DA, Finch BK, Asch SM. You are where you shop: grocery store locations,
378 weight, and neighborhoods. *American journal of preventive medicine*. 2006;31(1):10-17.
- 379 16. Liu GC, Wilson JS, Qi R, Ying J. Green neighborhoods, food retail and childhood overweight:
380 differences by population density. *American Journal of Health Promotion*. 2007;21(4s):317-325.
- 381 17. Leung CW, Laraia BA, Kelly M, et al. The influence of neighborhood food stores on change in
382 young girls' body mass index. *American journal of preventive medicine*. 2011;41(1):43-51.
- 383 18. Baker EA, Schootman M, Barnidge E, Kelly C. The role of race and poverty in access to foods that
384 enable individuals to adhere to dietary guidelines. *Prev Chronic Dis*. 2006;3(3):A76.
- 385 19. Barrera LH, Rothenberg SJ, Barquera S, Cifuentes E. The toxic food environment around
386 elementary schools and childhood obesity in Mexican cities. *American journal of preventive*
387 *medicine*. 2016;51(2):264-270.
- 388 20. Charreire H, Casey R, Salze P, et al. Measuring the food environment using geographical
389 information systems: a methodological review. *Public health nutrition*. 2010;13(11):1773-1785.
- 390 21. Moore LV, Roux AVD, Brines S. Comparing perception-based and geographic information system
391 (GIS)-based characterizations of the local food environment. *Journal of Urban Health*.
392 2008;85(2):206-216.
- 393 22. Moore LV, Diez Roux AV. Associations of neighborhood characteristics with the location and type
394 of food stores. *American journal of public health*. 2006;96(2):325-331.
- 395 23. Hernández B, Gortmaker SL, Laird NM, Colditz GA, Parra S, Peterson KE. Validez y
396 reproducibilidad de un cuestionario de actividad e inactividad física para escolares de la ciudad
397 de México. *Salud pública de México*. 2000;42(4).
- 398 24. Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: an update of
399 activity codes and MET intensities. *Medicine and science in sports and exercise*. 2000;32(9;
400 SUPP/1):S498-S504.
- 401 25. Lohman T, Roche A, Martorell R. *Standarization Reference Manual*. Champaign, IL: Human
402 Kinetics; 1988.
- 403 26. de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO
404 growth reference for school-aged children and adolescents. *Bull World Health Organ*.
405 2007;85(9):660-667.
- 406 27. Hill AM, LaForgia J, Coates AM, Buckley JD, Howe PR. Estimating abdominal adipose tissue with
407 DXA and anthropometry. *Obesity (Silver Spring, Md)*. 2007;15(2):504-510.
- 408 28. Ellis KJ, Abrams SA, Wong WW. Body composition of a young, multiethnic female population. *The*
409 *American journal of clinical nutrition*. 1997;65(3):724-731.
- 410 29. Ellis KJ. Body composition of a young, multiethnic, male population. *The American journal of*
411 *clinical nutrition*. 1997;66(6):1323-1331.
- 412 30. Chaudhari LS, Begay R, Schulz LO. Fifteen years of change in the food environment in a rural
413 Mexican community: the Maycoba project. *Rural Remote Health*. 2013;13(3):2404.

31. Monteiro CA, Levy RB, Claro RM, Castro IRRd, Cannon G. A new classification of foods based on the extent and purpose of their processing. *Cadernos de saude publica*. 2010;26:2039-2049.
32. Weaver CM, Dwyer J, Fulgoni III VL, et al. Processed foods: contributions to nutrition. *The American journal of clinical nutrition*. 2014;99(6):1525-1542.
33. Thornton LE, Pearce JR, Kavanagh AM. Using Geographic Information Systems (GIS) to assess the role of the built environment in influencing obesity: a glossary. *International Journal of Behavioral Nutrition and Physical Activity*. 2011;8(1):71.
34. Lind PL, Jensen PV, Glümer C, Toft U. The association between accessibility of local convenience stores and unhealthy diet. *The European Journal of Public Health*. 2016:ckv242.
35. Rose D, Richards R. Food store access and household fruit and vegetable use among participants in the US Food Stamp Program. *Public health nutrition*. 2004;7(08):1081-1088.
36. Daly CM, Foote SJ, Wadsworth DD. Physical Activity, Sedentary Behavior, Fruit and Vegetable Consumption and Access: What Influences Obesity in Rural Children? *Journal of community health*. 2017;42(5):968-973.
37. Larson NI, Story MT, Nelson MC. Neighborhood environments: disparities in access to healthy foods in the US. *American journal of preventive medicine*. 2009;36(1):74-81. e10.
38. Tang X, Ohri-Vachaspati P, Abbott JK, et al. Associations between food environment around schools and professionally measured weight status for middle and high school students. *Childhood Obesity*. 2014;10(6):511-517.
39. Cameron AJ, Thornton LE, McNaughton SA, Crawford D. Variation in supermarket exposure to energy-dense snack foods by socio-economic position. *Public health nutrition*. 2013;16(7):1178-1185.
40. Ghosh-Dastidar B, Cohen D, Hunter G, et al. Distance to store, food prices, and obesity in urban food deserts. *American journal of preventive medicine*. 2014;47(5):587-595.
41. Bodor JN, Rice JC, Farley TA, Swalm CM, Rose D. The association between obesity and urban food environments. *Journal of Urban Health*. 2010;87(5):771-781.
42. Cooksey-Stowers K, Schwartz M, Brownell K. Food swamps predict obesity rates better than food deserts in the United States. *International journal of environmental research and public health*. 2017;14(11):1366.
43. Lenardson JD, Hansen AY, Hartley D. Rural and remote food environments and obesity. *Current obesity reports*. 2015;4(1):46-53.
44. Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CA. The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Obesity*. 2015;23(7):1331-1344.
45. Zhou Q, Zhao L, Zhang L, et al. Neighborhood supermarket access and childhood obesity: a systematic review. *Obesity Reviews*. 2019.
46. Le H, Engler-Stringer R, Muhajarine N. Walkable home neighbourhood food environment and children's overweight and obesity: Proximity, density or price? *Canadian Journal of Public Health*. 2016;107(1):eS42-eS47.
47. Matanane L, Fialkowski MK, Silva J, et al. Para I Famagu'on-Ta: Fruit and Vegetable Intake, Food Store Environment, and Childhood Overweight/Obesity in the Children's Healthy Living Program on Guam. *Hawai'i Journal of Medicine & Public Health*. 2017;76(8):225.
48. Ellaway A, Macdonald L, Lamb K, Thornton L, Day P, Pearce J. Do obesity-promoting food environments cluster around socially disadvantaged schools in Glasgow, Scotland? *Health & place*. 2012;18(6):1335-1340.
49. Kestens Y, Daniel M. Social inequalities in food exposure around schools in an urban area. *American journal of preventive medicine*. 2010;39(1):33-40.

- 461 50. Rossen LM, Curriero FC, Cooley-Strickland M, Pollack KM. Food availability en route to school
462 and anthropometric change in urban children. *Journal of Urban Health*. 2013;90(4):653-666.
463 51. Strasburger VC. Children and TV advertising: nowhere to run, nowhere to hide. *Journal of*
464 *Developmental & Behavioral Pediatrics*. 2001;22(3):185-187.
465 52. Ashton D. Food advertising and childhood obesity. *Journal of the royal society of medicine*.
466 2004;97(2):51-52.
467 53. Caspi CE, Lenk K, Pelletier JE, et al. Association between store food environment and customer
468 purchases in small grocery stores, gas-marts, pharmacies and dollar stores. *The international*
469 *journal of behavioral nutrition and physical activity*. 2017;14(1):76.
470 54. Cameron AJ. The shelf space and strategic placement of healthy and discretionary foods in
471 urban, urban-fringe and rural/non-metropolitan Australian supermarkets. *Public Health Nutr*.
472 2018;21(3):593-600.

473

Table 1. Main characteristics of the study participants and according to body fat percentage (normal/high).

Characteristics	Total (n=218)		Normal body fat (n=105)		High body fat (n=113)	
	Mean ± SD ¹		Mean ± SD		Mean ± SD	
Age (months)	100.6 ±	18	97.14 ±	18.07	104 ±	18.34
Physical activity level						
Moderate physical activity (h/week)	8.57 ±	7.53	8.93 ±	8.06	8.24 ±	7.02
Intense physical activity (h/week)	3.37 ±	4.15	3.59 ±	5.00	3.15 ±	3.14
Anthropometry and body						
Weight (kg)	27.07 ±	8	23.36 ±	4.94	30.5 ±	8.4*
Height (cm)	125.3 ±	10	122.94 ±	9.28	127 ±	9.84*
Waist circumference (cm)	59.07 ±	9	54.12 ±	5.2	63.7 ±	9.25*
Height for age (z-score)	-0.72 ±	1	-0.89 ±	1.02	-0.56 ±	0.88
BMI ² for age (z-score)	0.33 ±	1	-0.34 ±	1.06	0.96 ±	1.02*
Body fat (%)	28.77 ±	7	23.42 ±	3.57	33.7 ±	4.78*
Abdominal fat (%)	27.82 ±	10	20.33 ±	4.32	34.8 ±	7.62*
	n	%	n	%	n	%
Caregiver's educational level						
No formal education	5	4.4%	14	6.4%	9	8.6%
Incomplete elementary school	28	24.8%	50	22.9%	22	21.0%
Complete elementary school	56	49.6%	114	52.3%	58	55.2%
Middle school or more	24	21.2%	40	18.3%	16	15.2%

¹ SD: Standard Deviation; ² BMI: Body Mass Index

*Different with t-test or Chi squared-test with Bonferroni correction for multiple comparisons between normal and high body fat (>30% girls and >25% boys).

482 **Table 2.** Linear meters of non or minimally and process foods from the convenience
 483 stores of the community

	Mean	SD ¹	Median	Minimum	Maximum
Non or Minimally Processed Foods	10.2	10.82	8.90	0	15.32
Processed Foods	16.72	9.32	16.78	0	22.57

¹ SD: Standard Deviation

484

485 **Table 3.** Density and proximity to convenience stores of the study participants according to body fat percentage
 486 (normal/high).

	Total (n=218)		Normal body fat (n=105)		High body fat (n=113)	
	Mean (SD) ¹	Median (range)	Mean (SD)	Median (range)	Mean (SD)	Median (range)
Density (n CS ² /250 m buffer)	11.5 (8.5)	11 (0 - 28)	9.6 (7.8)	10 (0-25)	12.8 (8.7) *	13 (0 - 28)
Proximity to closer store (m)	126.5 (107.2)	86.92 (5.5 - 683.8)	142 (106.7)	100.6 (5.8 - 378.5)	115.6 (107.0)	79.4 (5.6 - 682.8)

¹ SD: Standard Deviation; ² CS: Convenience stores

*Different with t-test with Bonferroni correction for multiple comparisons between normal and high body fat

487

Table 4. Association between food environment and measurements of obesity.

	Body fat %			Abdominal fat %			BMIz ³		
	β^1	95% CI ²	p^*	β	95% CI	p	β	95% CI	p
CS ⁴ Density (n/250 m buffer)	0.145	(0.048; 0.241)	0.004	0.206	(0.069; 0.343)	0.003	0.028	(0.005; 0.062)	0.005
Proximity (m)	-0.009	(-0.017; -0.001)	0.025	-0.012	(-0.023; -0.001)	0.033	-0.002	(-0.004; -0.001)	0.003
Food Access in CS (250m buffer)									
Shelf-space PF ⁵ (m)	0.071	(-0.249; 0.392)	0.662	0.047	(-0.409; 0.502)	0.845	0.002	(0.000; 0.003)	0.008
Shelf-space NMPF ⁶ (m)	0.055	(-0.105; 0.216)	0.497	0.049	(-0.178; 0.277)	0.661	0.072	(0.009; 0.135)	0.025
Ratio PF/NMPF	0.210	(-0.592; 0.479)	0.125	0.266	(-0.121; 0.654)	0.174	-0.012	(-0.061; 0.037)	0.662

¹ β : beta regression coefficient; ² BMIz: body mass index z-score; ³CI: 95% confidence interval; ⁴CS: convenience stores;

⁵PF: processed foods; ⁶NMPF: non or minimally processed foods

*Adjusted by: sex, age, caretaker's educational level, physical activity and proximity to school